

SHOT NOISE IN MESOSCOPIC QUANTUM SYSTEMS

M. P. Das

Department of Theoretical Physics
Research School of Physical Sciences and Engineering
The Australian National University
Canberra ACT 0200, Australia

F. Green

Centre for Quantum Computer Technology
School of Physics
University of New South Wales
Sydney, NSW 2052, Australia

1. INTRODUCTION

The fractional quantum Hall effect (FQHE) was discovered in nearly perfect two-dimensional electron systems, in the presence of a strong perpendicular magnetic field [1,2]. Such a field is sufficiently intense for the magnetic length to become comparable to the separation between carriers. Furthermore, the degenerate carriers (originally electrons and, later, holes) are quasiballistic. Their scattering mean free path is reckoned in microns.

The physics of current transport is particularly rich and exotic in the FQHE. Hence there is every reason to expect further puzzles – and a few surprises – in the fluctuation structure of this strongly correlated quantum Hall fluid. A particular class of theoretical tenets has come to dominate most, if not quite all, interpretations of noise measurements. From one point of view these particular notions could be seen as extending a relatively standard, semiclassical inspired treatment of fluctuations to quantum noise in the FQHE. In their own right, noise experiments for the FQHE are far from prosaic; they are remarkable *travaux de force* of experimental skill.

In these strongly quantized conductors, the longitudinal resistance displays remarkable oscillations as a function of the magnetic field B , dipping down to zero for finite intervals of B . At the same time, the transverse (Hall) resistance shows absolutely flat plateaux there. These features appear at specific, fractional values of the *filling factor*. The filling factor ν quantifies the state of the system: it is the ratio of the electron sheet density to the magnetic field (B can be expressed in units of flux quanta per unit area, thus counting the density of magnetic flux lines that pierce the electron sheet). When ν falls below one, we leave the domain of normal carrier behaviour and enter that of strong electron-electron correlations.

Below, we identify and critique some of the now-widespread beliefs about FQHE transport and fluctuations. We argue that these may not be as solidly grounded in the microscopics of the problem, as one would expect of a first-principles transport theory. This relative lack of well-defined grounding is already evident in certain semiclassical models of mesoscopic noise [3]. Similar questions of principle can be posed

in more highly quantum-coherent contexts; after all, essentially the same transport methodology is claimed to underlie both regimes (see, for example, Reference [4]). Current noise in the FQHE is a prime example for study. To set the scene, we first revisit the basic, low-field current response.

2. ESSENTIALS

The novel feature of Hall plateaux, observed at highly specific fractional fillings ($\nu = \frac{1}{3}, \frac{1}{5}$, and so on), was first explained by Laughlin [1] who devised an elegant many-body wave function to capture the behaviour of electrons in the FQHE state. It was soon realized that the elementary excitations of the FQHE must possess *fractional charge* and *fractional quantum statistics*. Following these discoveries, two issues emerged in the physics of the FQHE.

(1) At low applied voltage the Hall current is carried by the “edge states” located at the physical boundaries of the structure (see below). To probe this situation, a new transport experiment was set up to observe edge-state transport in a Corbino geometry [5]. Normally, in such a topology, carriers would necessarily transport current through the two-dimensional bulk. However, for any quantum Hall arrangement (both integral and fractional) the intricate potential landscape of the Landau levels, as sensed by the quasiparticles, should give rise to edge states. These would keep their one-dimensional character while meandering and threading their way right through the bulk of the Corbino disk.

The upshot of this and other experimental probes was that, for both integral and fractional quantum Hall situations, one came to believe that it is *only* the edge states that can channel the observable current. This leads to an unprecedented situation: topologically the edge states behave, to all intents, as strictly one-dimensional conductors [2] regardless of the two-dimensional nature of the original carrier states.

Contrary to the wide assumption that the edge states must carry all of the FQHE current in close neighborhood to the device boundary, von Klitzing has given a striking experimental counterexample. Careful Hall-conductance measurements in a Corbino structure show that, in his words, *the current cannot flow exclusively within a very narrow region close to the edge* [5].

(2) If a current is flowing, it should be possible to measure its associated *fluctuations* (noise). Noise exists in both equilibrium and nonequilibrium states of the system. From the scaling of the nonequilibrium fluctuations, and specifically those referred to as *shot noise*, one should be able to determine the effective charge quantum of the current-bearing excitations [4].

3. EDGE CHANNELS

Quantization of the Hall conductance in the integral (IQHE) and fractional quantum Hall effects is strikingly different in each case. This was pointed out by Beenakker and van Houten [6]. While a single-particle description (in the necessary presence of disorder) is invoked to explain the IQHE, the FQHE arises from many-body in-

teractions alone. Wherever the electronic Landau levels formed in the region of the sample boundaries intersect the Fermi level, they define a set of free current-carrying orbitals. These are known as the edge states. The boundaries are, from the carriers' point of view, regions of very high confining potential.

The response of carriers in the edge channels should depend directly on the form of the underlying Hamiltonian. Thus it matters critically whether the Landau-level scheme is an integral or a fractional one. In principle, experiments on edge-state conduction in the FQHE (strictly multi-particle physics) might well be expected to differ from the IQHE (strictly single-particle physics). Since, however, the current response appears rather similar in both, it is tempting to interpret *all* edge-state transport within a single, implicitly one-body, picture irrespective of whether the Landau-level filling is integral or fractional.

The assumption of a common mode for edge-state transport becomes much less self-evident in the case of the associated current fluctuations. Inherently, the latter will probe the many-body dynamical response even for a system of strictly *noninteracting* quantized fermions, free of short-range exchange-correlation effects. The manifest and fundamental distinction between the current, a *mean property*, and the noise, its mean-square statistical *variance*, goes to the heart of the most basic issues in the FQHE. While there is a systematic path from the latter to the former via the fluctuation-dissipation theorem, there is, of course, no general way of extracting the fluctuations purely from the mean current-voltage characteristics. While one may argue whether such issues have yet to be meaningfully addressed in mesoscopic physics, there is no doubt that they are basic to any understanding of the FQHE that claims to be physically complete.

In other words, current fluctuations are exquisitely sensitive to off-diagonal effects in a correlated system. They reveal much, much more than is contained in the current response by itself. The mean current response tells little – if anything – of the *uniqueness* of a multi-particle system's internal dynamics; the richness of its physics lies in the details of its off-diagonal behaviour. This more arduous analytical path is not usually the one that is trodden by popular accounts of FQHE noise.

The edge-state FQHE has been addressed simultaneously in three different theoretical papers [7,8,9], which provide three quite different answers to the problem. Fractional edge channels have also been studied in the laboratory by selective probing of the device boundaries [7,9]. In simple analogy with the IQHE, there has developed a picture in which FQHE edge channels carry the actual current via the *same* elementary modes that define the excitations of the FQHE deep in the bulk of the sample. A recognizably canonical, microscopic derivation of this intuitive hypothesis has not yet appeared in the literature.

Here one meets a delicate question: Just how is it that the edge excitations achieve the same internal configuration that is mandatory for the existence of the gapped, *incompressible* Laughlin quasiparticle states in the bulk? In our view, the theoretical constructs underpinning the present consensus on edge-state statistics – namely that they must be isomorphic with the excitations of the bulk FQHE – do not enjoy the clarity of reasoning evident in Laughlin's own theory of the incompressible bulk fractional states.

The appropriate formalism for the edge-state fractional excitations, and the ap-

appropriate experiments that would test it, are simply not in evidence so far. Such data as exists may (or may not) be suggestive of the intuitive consensus on the edge states' fractional inheritance from the Laughlin bulk. Unfortunately, we seem still to be far from a compelling proof that the consensus solution is unique, and that there is no alternative.

4. ONE-DIMENSIONAL CONDUCTION

Now we focus on a further assumption made for the edge states: the notion that they constitute a one-dimensional (1D) conductor very similar to the channel of a perfect 1D quantum wire. The only difference here is that edge excitations in the FQHE state are chiral. They move in only one direction around the boundary, and are physically decoupled from the bulk states (at the same time that they are supposed to inherit their fractionality from the Laughlin bulk).

The edge excitations in the IQHE are equivalent to a 1D noninteracting electron gas. The conductance G in this case is understood in terms of the quantized Landauer-Büttiker formula,

$$G \equiv \frac{I}{V} = 2 \frac{e^2}{h}. \quad (1)$$

(spin degeneracy yields the factor of two). In realistic cases, a subsequent transmission factor $\mathcal{T}$ is to be multiplied into the right hand side of Eq. (1) to account for nonideal forward scattering: if any scattering occurs in the channel, then $\mathcal{T} < 1$. For (chiral) edge states in either QHE, backscattering is automatically suppressed unless there are transitions that couple opposite edges (for instance, a constriction in the device). The Landauer-Büttiker formula seems to give a good description of edge-state transport in the idealized noninteracting problem [10].

For real systems, the electron-electron interaction cannot be ignored either. In 1D, even the weakest electron-electron interaction causes low-order perturbation theory to diverge, suggesting the breakdown of normal Fermi-liquid theory. One of two possible ways out of this breakdown is to carry out a careful dressing of the quasiparticles, either by resumming the most divergent diagrams or by introducing many-body effects through an effective exchange-correlation process. This should render the Fermi-liquid paradigm “almost” correct; that is, it becomes microscopically accurate in some asymptotic sense [11].

The other scenario is radically different. It proposes a model of a strongly localized, 1D interacting Fermi system known as the *Luttinger liquid* [12]. The Luttinger liquid has no relation to the perturbative Fermi-liquid picture. In this model, an infinitesimal interaction is enough to generate a multitude of (bosonic) collective modes. The excitation spectrum has absolutely no fermionic quasiparticle contributions in this case. An electronic state is filled by creation of an infinite number of bosons. This implies the separation of spin and charge degrees of freedom. Quite distinct excitations now carry spin and charge, which no longer coexist within the same state (as for a normal quasiparticle). The correlation functions of charge and spin are quite anomalous in comparison to those of a normal Fermi liquid. These are but a few of

many more differences between normal electrons and Luttinger excitations.

Measurements of transport in carefully fabricated quantum wires turn out to show departures from what is understood for the noninteracting conductance formula, Eq. (1). The universal conductance jump of e^2/h appears to be fractionalized. Therefore it seems natural to invoke the Luttinger-liquid model, which modifies the Landauer-Büttiker formula by a factor g that multiplies the right-hand expression of Eq. (1) [13]. It is to be noted that, formally, the same renormalized conductance can also arise from Kubo linear-response theory [14]. The value $g = 1$ is proper to noninteracting electrons. Any value other than unity will account for attractive ($g > 1$) or repulsive ($g < 1$) many-body interactions.

To explain the observed fractional conductance [13] by the Landauer-Büttiker formula, it is imaginatively proposed that the charge carriers responsible for transport become fractional and now carry the charge quantum $e^* = ge$, in keeping with Luttinger-liquid theory. We call the reader's attention to Ref. [14] for an interpretation of the role of the channel-lead contacts.

In the chiral fractional-charge model, the linear conductance is again $G \equiv I/V$. Whichever transport model is adopted, however, one has to renormalize *both* the current I and the operative voltage V (immediately across the source and drain boundaries of the channel). This holds for *all* theories of the interacting-electron liquid. It is then easy to show that, be it in a Fermi liquid or a Luttinger liquid, the current and voltage are renormalized to precisely the same extent. As a result, the conductance given in Eq. (1) stays exactly as one finds it in a noninteracting system.

In the Luttinger-liquid interpretation, the 1D quantum wire is purely ballistic, free from impurities and thus collisionless. Its resistance can come only from coupling of the channel's contacts with the macroscopic, and dissipative, outer source and drain leads. The disordered leads are normal metallic Fermi liquids, not 1D systems. Their large number of internal degrees of freedom provides the excitations that engender the so-called "contact" resistance. The properties of the contacts make no reference to details of the excitations that dominate within the channel itself. Indeed, this conceptual division between the physical channel and the physical leads is essential to the applicability of the Landauer-Büttiker conductance [15]. However, the exact role of disorder is yet to be spelled out microscopically within this picture. One has only the assertion that the mismatch between densities of states in channel and leads, is a self-sufficient explanation.

The actual nature of the contact-resistance problem is far from trivial. It has been considered in some depth by Fenton [16] among others, and more recently by Magnus and Schoenmaker [17]. Even granting the consensus view on the issue of the densities of states and their mismatch, this still leaves one to work out a theory for the intrinsic states of the channel itself. Let us look at that.

In a provocative theoretical paper [18] Wen first argued that in the FQHE situation (at filling $\nu < 1$ such that ν^{-1} is an odd integer) the edge-current modes are isomorphic to the Luttinger-liquid modes. In a quantum-Hall-bar geometry the right- and left-moving excitations are localized near the top and bottom of the bar, respectively. The chemical-potential difference between the top and bottom edge excitations is identified with the drop in the Hall potential, normal to the current. The Hall conductance is obtained as $G = ge^2/h$. Here $g = \nu^{-1}$ is nothing other

than the odd-integer fractional filling. One now considers the situation for inter-edge tunnelling at a point contact: the tunneling current between the top and bottom edge states is claimed to be transported physically by charges in units of $e^* = e/\nu$.

Another intriguing aspect of the edge-tunneling experiments is nonlinear scaling of the I - V characteristic. The power-law exponent in the predicted relation $I \sim V^g$ shows a direct proportionality to the inverse of the filling factor, as expected within the Luttinger picture. As the fractional filling changes, the exponent should also change, discontinuously. This is *not* what is observed in the laboratory [19].

At this point it suffices to remark that the theory of edge construction is yet to be clearly articulated. Two contrasting points of view are advanced by Beenakker [7] and MacDonald [8]. MacDonald argued that, if the edge width is smaller than the magnetic length, the edge channel would be very sharp, indeed abrupt enough to support a strictly 1D Luttinger liquid. One could then have the Laughlin excitations travelling about the edge and acting as Luttinger excitations. Against this argument, Beenakker pointed out that a realistic edge width will be appreciably greater than the magnetic length; in that case, the edge current must be carried by normal electrons rather than fractionalized charge states. He further argues that, for intermediate edge widths, one might observe fractional filling factors whose values are different from the bulk.

Experiments on shot noise in FQHE devices have not been supplemented by measurements of the actual edge width in their structures. On the other hand, groups who have measured the quantized conductance appear to see edge widths that are smooth, rather than abrupt, on the magnetic-length scale. The edges seem to support complicated multiple-channel structures, consistent with Beenakker's account [20].

5. FLUCTUATIONS

Shot noise in electronic conduction is a consequence of the *granularity* of the electronic charge. If successive arrival times for discrete charge carriers at the collector electrode have a Poissonian distribution, one obtains the spectral distribution for shot noise $S(0) = 2q\bar{I}$. Here the spectral function $S(0)$ is the zero-frequency Fourier transform of the current-current correlation function; q is the charge quantum of the current carriers and $\bar{I}$ is the time-averaged current. The above formula for (classical) shot noise is a famous result, known in the literature as the Schottky formula. It is clearly a signature of *nonequilibrium* charge dynamics in the conducting system. Shot noise exhibits no dependence on temperature, quite unlike thermally generated noise, whose current fluctuations at finite temperature T , and low voltage $eV \ll k_B T$ (the quasi-equilibrium limit), yield the Johnson-Nyquist thermal spectrum $S_{\text{JN}}(0) = 4Gk_B T$. This typically scales both with temperature and conductance.

Currently, the dominant theories of electronic conduction and shot noise in mesoscopic systems all rely on a concept of *independent* quasiparticle motion, namely the Landauer-Büttiker formalism for coherent single-carrier transmission. Such a description has shown enormous flexibility in applications to transport in multifarious physical systems [4]. At the same time, it pays little heed to any role for interactions among the carriers of a mesoscopic system. In the large majority of mesoscopic

studies that have proliferated during the past decade, many-body interactions are assumed to play a minor part in the scheme of things [15]. It is widely supposed that – if they matter at all – the interactions can be included as a kind of perturbative afterthought.

Below we look, qualitatively, at just a few of the reasons why consideration of the interactions is by no means an optional extra. Indeed, many-body interactions lie at the very heart of fluctuation kinetics. Any model failing to account for them properly, especially in degenerate Fermi liquids, risks serious inconsistency with the conservation laws and the sum rules [3]. Without a guarantee of microscopic conservation, there can be no assurance that one’s predictions make physical sense.

6. SHOT-NOISE EXPERIMENTS

Two experimental groups, Saminadayar *et al.* [21] at Saclay and de Picciotto *et al.* [22] at the Weizmann Institute, have reported the direct observation of fractional charges in FQHE systems. These works measured the shot-noise spectral function for the current. Measurements were done on the edge states of a high-mobility two-dimensional electron gas, within a quantum point contact. This geometry was created by controlling the bias voltage on a split gate, constricting the 2DEG locally and bringing the edge states close enough for tunnelling to occur. Filling factors, either $\frac{1}{3}$ or $\frac{1}{5}$, were fixed at will by modulating the areal densities of the electrons and the magnetic flux.

In this way shot noise could be measured at very low frequencies and at very low temperatures (in the range of mK). The current cross-correlation (essentially shot noise) was found to be proportional to the back-scattered current $I_B = I_0 - \bar{I}$, where $\bar{I}$ is the mean transmitted current and $I_0 = (e^*/h)eV$ is the (chiral) current coming into the quantum point contact. Most importantly, e^* is the elementary quasiparticle charge ($e/3$, $e/5$, etc.). The results from both Saclay and Weizmann generally show that the shot noise scales nicely with the back-scattered current. The best-fit slope scales with e^* , giving evidence for fractional charge quantization.

On closer comparison of the two experimental accounts, one finds some significant differences in data interpretation. Here we recall that, according to Landauer-Büttiker theory, the Schottky formula requires correction by the Fano factor γ which, if less than unity, suppresses the spectral density so that $S(0) = 2\gamma qI$. In any sort of barrier, such as a quantum point contact, this suppression is governed solely by the transmission coefficient $\mathcal{T} \leq 1$. For a single conducting channel, the backscattered shot noise should carry the Fano factor $\gamma = \mathcal{T}$.

The data of de Picciotto *et al.* [22] was phenomenologically fitted by choosing a certain value of $\mathcal{T}$ to improve agreement with predictions for backscattering; then $S(0) \rightarrow 2e^*I_B\mathcal{T}$. Saminadayar *et al.* [21], on the other hand, fitted their backscattering data without *any* suppression factor at all, rightly pointing out that the presence of a suppression factor would mean that the actual quasiparticle charge must exceed its fundamental value (in their case, $e/3$), to offset the action of $\mathcal{T}$ and recover the linear $e/3$ slope as observed by them. Yet, despite these mutual inconsistencies it is nevertheless asserted that the fractional charge of the (bulk) Laughlin quasiparticles

has indeed been measured, and in the most direct way possible, by these shot-noise experiments [23].

7. THEORY

Now we ask the question: What is the basis for understanding the above experiments? Let us start with a pure, low-density, interacting 2D electron gas in a high magnetic field.

The system has degenerate Landau levels. As shown by Laughlin [1] the quasiparticles in this system at fractional filling $\nu = 1/q$ (q odd) have fractional charge $e^* = e/q$ as well as an energy gap $D = e^2/l$ where $l = \sqrt{\hbar/eB}$ is the Landau length. The quasiparticle is an electron with q flux quanta associated with it.

For a finite system, the flat Landau levels of the bulk bend upward in energy owing to the confining potential. Crossing of the Fermi level with the confining potential creates the edge states. For the IQHE the edge-state quasiparticles are electrons with charge e . For the FQHE, by analogy, the edge-state quasiparticles are taken to be the Laughlin excitations of the bulk, with charge e^* . Now the edge channels behave like 1D chiral Luttinger liquid.

Theories of quantum noise due to the Luttinger excitations are available in Refs. [24-27]. Here, Luttinger excitations of the interacting 1D electrons and Laughlin excitations of the bulk FQHE are regarded as synonymous. Kane and Fisher [24] treated the edge excitations as a Luttinger liquid and used a bosonization technique. There are two tunneling regimes: strong and weak.

Strong tunneling leads to weak backscattering, and vice versa. The experiments with quantum point contacts, described in the Section above, relate to the weak-tunneling case, with strong backscattering. The formula for shot noise is of precisely the same form as the noninteracting Landauer-Büttiker formula. Chamon *et al.* [25] have discussed the strong and weak tunneling limit of Luttinger excitations and used a nonequilibrium Keldysh formalism to obtain nonlinear current-voltage relationship, as expected for a Luttinger liquid. Fendley *et al.* [26] have considered a conformal field-theoretic approach to obtain the shot noise of Luttinger liquids, and have obtained strong- and weak-tunneling results at finite temperature.

In a more recent paper Sandler *et al.* [27] have considered tunneling between integral states and various fractional states. Here the fractionally charged states are not necessarily Laughlin quasiparticles. Rather, they correspond to solutions of the coupled systems. Generalizations of these results to strong- and weak-coupling regimes are made for a quantum-point-contact geometry.

8. SUMMARY AND CONCLUSIONS

With a 1D Luttinger liquid as their starting point, all of these various theories ultimately converge to the standard Landauer-Büttiker formalism. This has exhibited great success in understanding a huge variety of physical systems [3]. However, for the measurement and interpretation of shot-noise results, it is unclear how these

approaches, in conjunction with the Landauer-Büttiker picture, manage to retain control of the fluctuation structure, whose off-diagonal properties manifest the intrinsically *correlated* nature of such strongly interacting systems.

We believe that the following points warrant more complete and logically coherent explanations than are available at this writing.

- For the fractional edge-channel state, commonly considered as a Luttinger liquid, one needs to have a theory with truly clear, physically well-formed premises.
- Why should the fractional charge of the Luttinger liquid be at all the same as the fractional charge of the bulk Laughlin excitations?
- How do the bulk, gapped, incompressible excitations lose their gap and their incompressible character in going to the boundary?
- The quantum shot-noise formula has the typical Schottky form, corrected for the Landauer-Büttiker suppression in forward transmission, $1 - \mathcal{T}$ (or $\mathcal{T}$ for backscattering). True many-body statistics and correlations are totally missing in this one-body formula. Recently Isakov *et al.* [28] tried to incorporate exclusion statistics, but at the cost of even greater conceptual difficulties that undermine the heuristic state-counting argument [4,29].

The important issues related to shot noise have been covered recently in the extensive review by Blanter and Büttiker [4]. While transport and noise are considered mainly in the context of independent quasiparticles, those authors themselves state that “electrons are, however, interacting entities and both the fluctuations at finite frequencies and the fluctuation properties far from equilibrium require in general a discussion of the role of the long range Coulomb interaction. A quasi-particle picture is no longer sufficient and collective properties of the electron system come into play.”

To correctly understand both conductance and shot-noise experiments in a unified way, one has to develop a genuinely nonequilibrium theory of transport and fluctuations, including the electron correlations that have so far been neglected (or, at best, averaged over in an *ad hoc* fashion). The Landauer-Büttiker formula is derivable from Kubo linear-response theory, and this provides at least one formal link back to the many-particle density matrix and its inbuilt correlations. While recognizing the successes of Landauer-Büttiker theory, we feel that the time is now ripe to formulate a practical and correct *many-body* theory for the complex behavior of correlated mesoscopic systems.

REFERENCES

- [1] See *Nobel Lectures: Fractional Quantum Hall Effect*, Rev. Mod. Phys. **71**, 4 (1998)
- [2] T. Chakraborty and P. Pietilainen, *The Fractional Quantum Hall Effect* (Springer, Berlin, 1988).
- [3] F. Green and M. P. Das, *J. Phys: Condens. Matter* **12**, 5233 (2000).
- [4] Ya. M. Blanter and M. Büttiker, *Phys. Rep.* **336**, 1 (2000).
- [5] K. von Klitzing, *Physica B* **184**, 1 (1993).
- [6] C. W. J. Beenakker and H. van Houten in *Advances in Solid State Physics*, Vol.

- 44, edited by H. Ehrenreich and D. Turnbull, (Academic, New York, 1991).
- [7] C. W. J. Beenakker, *Phys. Rev. Lett.* **64**, 216 (1990).
 - [8] A. H. MacDonald, *Phys. Rev. Lett.* **64**, 220 (1990).
 - [9] A. M. Chang, *Solid State Comm.* **74**, 871 (1990).
 - [10] B. J. van Wees, H. van Houten, C. W. J. Beenakker, J. G. Williamson, L. P. Kouwenhoven, D. J. van der Marel, and C. T. Foxon, *Phys. Rev. Lett.* **60**, 848 (1988); D. A. Wharam, M. Pepper, H. Ahmed, J. E. F. Frost, D. G. Hasko, D. C. Peacock, D. A. Ritchie, and G. A. C. Jones, *J. Phys. C* **21**, L209 (1988).
 - [11] See, for example, S. Das Sarma and B. Hu, *Aust. J. Phys.* **46**, 359 (1993).
 - [12] G. D. Mahan, *Many Particle Physics* (Plenum, New York, 1990).
 - [13] A. Yacoby, H. L. Störmer, N. S. Wingreen, L. N. Pfeiffer, K. W. Baldwin, and K. W. West, *Phys. Rev. Lett.* **77**, 4612 (1996); K. J. Thomas, J. T. Nicholls, N. J. Appleyard, M. Y. Simmons, M. Pepper, D. R. Mace, W. R. Tribe, and D. A. Ritchie, *Phys. Rev. Lett.* **77**, 135 (1996).
 - [14] C. Kane and M. P. A. Fisher, *Phys. Rev. B* **46**, 15233 (1992); M. P. A. Fisher and L. I. Glazman in *Mesoscopic Electron Transport (NATO ASI Series E)* ed L. P. Kouwenhoven, G. Schön, and L. L. Sohn (Kluwer Academic, Dordrecht, 1997).
 - [15] Y. Imry and R. Landauer, *Rev. Mod. Phys.* **71**, S306 (1999).
 - [16] E. W. Fenton, *Phys. Rev. B* **46**, 3754 (1992); *Superlattices and Microstruct.* **16**, 87 (1994).
 - [17] W. Magnus and W. Schoenmaker, *Phys. Rev. B* **61**, 10883 (2000).
 - [18] X.-G. Wen, *Phys. Rev. Lett.* **64**, 2206 (1990); *International J. Mod. Phys. B* **6**, 1711 (1992).
 - [19] M. Grayson, D. C. Tsui, L. N. Pfeiffer, K. W. West, and A. M. Chang *Phys. Rev. Lett.* **80**, 1062 (1998).
 - [20] M. Ando, A. Endo, S. Katsumoto, and Y. Iye, *Physica B* **249-251** (1998), pp 426-9.
 - [21] L. Saminadayar, D.C. Glattli, Y. Jin, and B. Etienne, *Phys. Rev. Lett.* **79**, 2526 (1997); D. C. Glattli, V. Rodriguez, H. Perrin, P. Roche, Y. Jin, and B. Etienne, *Physica E* **6**, 22 (2000).
 - [22] R. de Picciotto, M. Reznikov, M. Heiblum, V. Umansky, G. Bumin, and D. Mahalu, *Nature* **389**, 162 (1997); M. Reznikov, R. de Picciotto, T. G. Griffiths, M. Heiblum, and V. Umansky, *Nature* **399**, 238 (1999).
 - [23] M. Heiblum and A. Stern, *Physics World* **13**, 37 (2000).
 - [24] C. L. Kane and M. P. A. Fisher, *Phys. Rev. Lett.* **72**, 724 (1994).
 - [25] P. Fendley, A. W. W. Ludwig, and H. Saleur, *Phys. Rev. Lett.* **75**, 2196 (1995).
 - [26] C. de C. Chamon, D. E. Freed, and X. G. Wen, *Phys. Rev. B* **51**, 2363 (1995).
 - [27] N. P. Sandler, C. de C. Chamon, and E. Fradkin, *Phys. Rev. B* **59**, 12521 (1999).
 - [28] S. B. Isakov, Th. Martin, and S. Ouvry. *Phys. Rev. Lett.* **83**, 580 (1999).
 - [29] M. P. Das and F. Green, *Phys. Rev. Lett.* **85**, 222 (2000).